

LETTER TO JMG

Mutations of the Birt-Hogg-Dubé (*BHD*) gene in sporadic colorectal carcinomas and colorectal carcinoma cell lines with microsatellite instability

J-H Shin, Y-K Shin, J-L Ku, S-Y Jeong, S-H Hong, S-Y Park, W-H Kim, J-G Park

J Med Genet 2003;40:364–367

Birt-Hogg-Dubé (*BHD*) syndrome, an inherited autosomal genodermatosis characterised by benign tumours of the hair follicle, is associated with renal neoplasia, lung cysts, and spontaneous pneumothorax.¹ The novel causative gene, identified by linkage analysis in *BHD* families, is localised on chromosome 17p11.2.² Protein truncating germinal mutations within a hypermutable (C)₈ tract occur in patients with *BHD* syndrome and lead to an increased risk of kidney cancer.³

Microsatellite repeats are widely distributed throughout the genome. Owing to a defect in the DNA mismatch repair gene, a subset of tumours accumulates frequent deletion and insertion mutations in these repetitive DNA sequences.^{4–6} Most microsatellite instability (MSI) has so far been described in non-coding DNA within introns of intergenic regions in the genome. However, in some cancer related genes and mismatch repair genes, MSI has been identified in protein coding regions. The first target sequence identified within a coding region was a poly (A)₁₀ nucleotide tract of the *TGFBR2* gene.⁷ The other mutational targets of MSI have been found in repetitive sequences of *IGF2R*⁸ and *BAX*⁹ genes involved in the regulation of cell growth and in the promotion of apoptosis, respectively. Furthermore, frameshift mutations in repeat sequences of the DNA mismatch repair genes *MSH3* and *MSH6* have been reported.¹⁰ In tumours with MSI, the mechanism of tumorigenesis is believed to involve frameshift mutations of microsatellite repeats within the coding regions of genes, the inactivation of which is considered to contribute to tumorigenesis.

Early reports suggested that *BHD* syndrome was associated with a predisposition to colon neoplasms,^{11–13} but Zbar *et al*¹⁴ reported that colon cancer and colon polyps are not related to *BHD* syndrome. Recently, Khoo *et al*¹⁵ reported that colorectal neoplasia is an associated feature of *BHD* in some families. To elucidate whether the *BHD* gene is associated with colon neoplasia and one of the MSI target genes, we screened the poly (C)₈ tract of the *BHD* gene, a mutational hot spot, in 32 MSI sporadic colorectal carcinomas and 80 microsatellite stable (MSS) sporadic colorectal carcinomas. In addition, we screened the entire coding region of the *BHD* gene in 13 MSI colorectal carcinoma cell lines and nine MSS colorectal carcinoma cell lines.

MATERIALS AND METHODS

Samples

As described in our previous study,¹⁶ we collected 325 consecutive patients who had undergone operation for primary colorectal cancers at Seoul National University Hospital, Seoul, Korea. Fifteen cases of familial colorectal cancers (two HNPCC, four FAP, and nine with a family history of cancer) were excluded through detailed family history taking, obtained by interviewing the patients and their relatives. Twenty-two colorectal cancer cell lines were obtained from the Korean Cell Line Bank, Seoul, Korea. Thirteen of these cell

Key points

- Birt-Hogg-Dubé (*BHD*) syndrome, an inherited autosomal genodermatosis characterised by benign tumours of the hair follicle, is associated with renal neoplasia, lung cysts, and spontaneous pneumothorax.
- Recently, the *BHD* gene containing a potential MSI target sequence has been identified as a cancer causing gene, which encodes a novel protein called folliculin.
- Although earlier studies described the relation between colonic neoplasia and *BHD* syndrome, it was questionable that the *BHD* gene was associated with colon neoplasia.
- We screened the poly (C)₈ tract of the *BHD* gene, a mutational hot spot, and found mutations in five (16%) of 32 MSI sporadic colorectal carcinomas and in one (7.7%) of 13 MSI colorectal carcinoma cell lines. However, we were unable to find any frameshift mutation in 80 MSS sporadic carcinomas or nine MSS colorectal carcinoma cell lines. In addition, we found two heterozygous missense mutations, Arg137Cys and Arg462Ser, in SNU-1040 and LoVo cell lines with MSI, respectively.
- The frequency (16%) of mutations in the *BHD* gene was the same as that in the *IGF2R* gene and this was less than that of the *TGFBR2*, *MSH3*, *BAX*, and *MSH6* genes. Interestingly, all tumours with the *BHD* gene mutation harboured concurrent mutations of the poly (C)₈ tract of the *MSH6* gene ($p=0.002$), but the frameshift mutations of the *BHD* and *IGF2R* genes were mutually exclusive.
- Our findings strongly support that the *BHD* gene is associated with colon cancer and putative MSI target genes involved in the development of MSI colorectal carcinomas.

lines (SNU-175, SNU-407, SNU-769A, SNU-769B, SNU-1040, SNU-1047, SNU-C2A, SNU-C4, HCT-116, LoVo, LS174T, HCT-8, and DLD-1) are MMR negative and nine (SNU-61, SNU-81, SNU-283, SNU-503, SNU-1033, SNU-1197, Colo205, SW480, and WiDr) are MMR positive in terms of their MSI status, as determined by a previous study.^{17–19}

DNA isolation and MSI analysis

In a total of 310 cases of sporadic colorectal cancers, genomic DNA from the tumour and corresponding normal tissues were procured from formalin fixed and paraffin embedded tissue samples by microdissection. MSI analysis of tumours was performed using an ABI 377 automatic sequencer (Perkin-Elmer,




Figure 1 Sequencing analysis of the representative frameshift mutations of the *BHD* gene in MSI positive sporadic colorectal carcinoma. Arrows indicate insertion/deletion sequences in mononucleotide repeats. The wild type poly (C)₈ tract is marked in bold letters.

Foster City, CA) with fluorescent dye labelled primers of BAT-26 and BAT-25 markers. Thirty-two of 310 (10.3%) sporadic colorectal carcinomas were defined as MSI.

Mutation analysis of the *BHD* gene

To screen for mutations in the *BHD* gene, entire exons of the gene were examined by DNA sequence analysis. The primer sequences and the detailed reaction conditions for amplification have been described previously.³ Bidirectional sequences of PCR products were analysed using an ABI Prism 3100 Genetic Analyzer (Applied Biosystems, Foster City, CA).

Statistical analysis

To compare the frequencies of frameshift mutations of MSI target genes, we performed the χ^2 test or Fisher's exact test using SPSS software (version 10.05, SPSS Inc, Illinois, USA). Odds ratio (OR) and 95% confidence intervals (CI) were also calculated. A p level of <0.05 was considered statistically significant.

RESULTS

Three different mutations, a 1 bp and 2 bp deletion and a 1 bp insertion in the (C)₈ repeat of the coding region of the *BHD* gene, were observed in five (16%) of 32 sporadic colorectal carcinomas (fig 1). In addition, HCT-116 in the MSI cell lines had a 1 bp insertion mutation in the poly (C)₈ tract (1 of 13, 7.7%). As controls, none of 80 MSS sporadic colorectal carcinomas or the nine MSS colorectal carcinoma cell lines had mutations in the (C)₈ repeat of the coding region of the *BHD* gene. The overall mutational profiles of the six genes evaluated showed diverse combinations (table 1). Among the 32 MSI tumours, 31 (97%) tumours had mutations in more than one gene, 25 (78%) tumours had mutations in more than two genes, 15 (47%) tumours had mutations in more than three genes, and five (16%) tumours had mutations in more than four genes. There were no homozygous mutations in the *BHD*, *IGF2R*, or *MSH3* genes. Homozygous mutations in the *BAX* and *MSH6* genes were rare, only one of 16 *BAX* mutations and two of 11 *MSH6* mutation, whereas most of the *TGFBR2* mutations were homozygous (13 out of 26, 50%). The frequency (16%) of mutations in the *BHD* gene was the same as that in the *IGF2R* gene and this was less than that of the *TGFBR2*, *MSH3*, *BAX*, and *MSH6* genes. Interestingly, although the expected ratio of concomitant *BHD-MSH6* mutation was only 5.4%, all tumours with the *BHD* gene mutation harboured

concurrent mutations of the poly (C)₈ tract of the *MSH6* gene (p=0.002, OR =1.833, 95% confidence intervals 1.069 to -3.144), but the frameshift mutations of the *BHD* and *IGF2R* genes were mutually exclusive. Like tumours with a *BHD* frameshift mutation, HCT-116 harboured concurrent mutation of *MSH6* and not *IGF2R*.

To investigate further whether other types of mutation were found in both MSI and MSS colorectal cancer cell lines, we screened the entire coding region of *BHD* in 13 MSI cell lines and nine MSS cell lines. Two heterozygous missense mutations, Arg137Cys and Arg462Ser, were found in SNU-1040 and LoVo cell lines with MSI, respectively. However, no biallelic mutation was found in 13 MSI and nine MSS cell lines.

DISCUSSION

BHD syndrome is a rare inherited genodermatosis characterised by hair follicle hamartomas, kidney tumours, and spontaneous pneumothorax. Protein truncating mutations have been found in the *BHD* gene in BHD families, and most reported insertion/deletion mutations have been found within the poly (C)₈ tract.^{3 15} These results suggest that the polycytosine mononucleotide tract is hypermutable and particularly prone to disease causing mutations. These insertion/deletion mutations resulted in a stop codon predicted to truncate the protein 26 missense amino acids downstream and 38 missense amino acids downstream, respectively. Truncated proteins lose the *N*-glycosylation site and the myristoylation site.³ In addition, we found two heterozygous missense mutations, Arg137Cys and Arg462Ser, in SNU-1040 and LoVo cell lines with MSI, respectively. These missense mutations result in an amino acid change from positively charged to uncharged amino acid. In addition, BHD protein, folliculin, was highly conserved among mouse, *D melanogaster*, and *C elegans* and this novel protein may have an important biological function in a wide range of organisms. Clues to the mechanism of tumorigenesis in sporadic colorectal carcinoma with a defective *BHD* gene may come from consideration of the BHD syndrome phenotype. Phenotypic data¹ suggest that defective folliculin may affect the structure of the cytoskeletal network and the interaction between epithelial and mesenchymal cells. Perhaps the mutation alters the composition of the extracellular matrix, and produces a structural or microenvironmental abnormality that affects the regulation of cellular proliferation.

Table 1 Mutation profile of the MSI target genes in 32 MSI positive sporadic colorectal carcinomas

Case No	Target genes					
	TGFBR2 (A) ₁₀	BAX (G) ₈	IGF2R (G) ₈	MSH3 (A) ₈	MSH6 (C) ₈	BHD (C) ₈
24	-2/-2	wt	wt	-1/wt	wt	wt
45	wt	wt	wt	wt	-1/wt	-2/wt
51	wt	wt	wt	wt	-1/wt	wt
59	-1/+1	wt	wt	wt	-1/wt	+1/wt
66	-1/-2	wt	wt	wt	-1/wt	+1/wt
67	-1/-2	-1/-1	wt	-1/wt	wt	wt
68	-1/+1	wt	wt	wt	wt	wt
90	-1/wt	wt	wt	-1/wt	-1/wt	wt
91	-1/-1	wt	wt	-1/wt	-1/wt	-1/wt
95	-2/wt	-1/wt	wt	-1/wt	-1/-1	-1/wt
102	-2/+1	wt	-1/wt	-1/wt	-1/-1	wt
134	wt	-1/wt	wt	wt	wt	wt
151	-1/-2	-1/wt	wt	wt	wt	wt
153	-1/-2	-1/wt	wt	wt	-1/wt	wt
164	-1/wt	-1/wt	-1/wt	-1/wt	wt	wt
171	-1/-1	-1/wt	wt	-1/wt	wt	wt
172	-1/-2	-1/wt	+1/wt	-1/wt	wt	wt
193	-1/wt	-1/wt	wt	wt	wt	wt
206	-1/wt	-1/wt	-1/wt	wt	wt	wt
222	-2/-1	-1/wt	wt	-1/wt	wt	wt
218	wt	wt	wt	wt	wt	wt
225	-1/wt	wt	wt	-1/wt	wt	wt
236	-1/wt	wt	-1/wt	wt	wt	wt
243	-2/wt	-1/wt	wt	wt	wt	wt
256	-1/-2	-1/wt	wt	wt	-1/wt	wt
260	wt	wt	wt	-1/wt	wt	wt
268	-1/wt	wt	wt	wt	wt	wt
380	wt	-1/wt	wt	-1/wt	wt	wt
381	-1/-1	wt	wt	wt	wt	wt
408	-1/wt	-1/wt	wt	-1/wt	wt	wt
411	-1/wt	wt	wt	wt	-1/wt	wt
444	-2/wt	-1/wt	wt	wt	wt	wt
	26 (81%)	16 (50%)	5 (16%)	14 (44%)	11 (34%)	5 (16%)

Previously, Shin *et al*⁶ reported the status of mutations in five known target genes. wt, wild type. -1/+1, 1 base deletion/1 base insertion. Bold letters = tumours identified as MSI in this study.

In most published studies of potential target genes, the prevalence of SMT (short mononucleotide tract) mutations in control, presumably unselected genes has been noted to be low. The study of such control genes has in general indicated that A or G tracts of 9 bp or fewer are mutated in <5% of cancers with MSI. Duval *et al*²⁰ proposed an experimental cut off frequency value (10-15%) that divides mutational events into those that are or are not selected for during MSI-H tumour progression. In a recent paper,²¹ although real target genes were confirmed by comparing the frequency of mutation in the coding repeat with the non-coding repeat, these results were dependent on different comparative statistical approaches and showed the difference of significance. The mutation frequency is still likely to be a useful indicator of the functional role played by a specific gene. According to our results that 16% of MSI tumours harboured mutations of the poly (C)₈ tract in the *BHD* gene, there is still reason for concluding that the *BHD* gene is a putative MSI target gene. To confirm that the *BHD* gene is a target of MSI colorectal cancer, general cytological approaches using gene array technologies are needed.

In summary, we found that mutations of the poly (C)₈ tract of the *BHD* gene in MSI sporadic colorectal carcinomas and the frequency of mutations in the *BHD* gene is comparable with that of the *IGF2R* gene, which is one of the well known MSI target genes. In addition, two heterozygous missense mutations, Arg137Cys and Arg462Ser, were found in SNU-1040 and LoVo cell lines with MSI, respectively. Functional studies are required to understand the biological role of follinulin in colorectal carcinogenesis. Our findings suggest that the *BHD* gene is associated with colon cancer and putative MSI target genes involved in the development of MSI colorectal carcinomas.

ACKNOWLEDGEMENTS

This work was supported by the National Cancer Centre Research Grant (Research Grant 2001) and the 2002 BK21 project for Medicine, Dentistry, and Pharmacy.

Authors' affiliations

J-H Shin, Y-K Shin, J-L Ku, S-H Hong, J-G Park, Laboratory of Cell Biology, Cancer Research Institute and Cancer Research Centre, Seoul National University College of Medicine, Seoul 110-744, Korea
S-Y Jeong, S-Y Park, J-G Park, Research Institute and Hospital, National Cancer Centre, Goyang, Gyeonggi 411-764, Korea
W-H Kim, Department of Pathology, Seoul National University College of Medicine, Seoul 110-744, Korea

Correspondence to: Dr J-G Park, National Cancer Centre, 809 Madu-dong, Ilsan-gu, Goyang, Gyeonggi 411-764, Korea; park@ncc.re.kr

REFERENCES

- Schmidt LS**, Warren MB, Nickerson ML, Weirich G, Matrosova V, Toro JR, Turner ML, Duray P, Merino M, Hewitt S, Pavlovich CP, Glenn G, Greenberg CR, Linehan WM, Zbar B. Birt-Hogg-Dubé syndrome, a genodermatosis associated with spontaneous pneumothorax and kidney neoplasia, maps to chromosome 17p11.2. *Am J Hum Genet* 2001;**69**:876-82.
- Khoo SK**, Bradley M, Wong FK, Hedblad MA, Nordenskjöld M, The BT. Birt-Hogg-Dubé syndrome: mapping of a novel hereditary neoplasia gene to chromosome 17p12-q11.2. *Oncogene* 2001;**23**:5239-42.
- Nickerson ML**, Warren MB, Toro JR, Matrosova V, Glenn G, Turner ML, Duray P, Merino M, Choyke P, Pavlovich CP, Sharma N, Walther M, Munroe D, Hill R, Maher E, Greenberg C, Lerman MI, Linehan WM, Zbar B, Schmidt LS. Mutations in a novel gene lead to kidney tumors, lung wall defects, and benign tumors of the hair follicle in patients with the Birt-Hogg-Dubé syndrome. *Cancer Cell* 2002;**2**:157-64.
- Aaltonen LA**, Peltomäki P, Leach FS, Sistonen P, Pylkkänen L, Mecklin JP, Järvinen H, Powell SM, Jen J, Hamilton SR, Petersen GM, Kinzler KW, Vogelstein B, de la Chapelle A. Clues to the pathogenesis of familial colorectal cancer. *Science* 1993;**260**:812-16.

- 5 **Ionov Y**, Peinado MA, Malkhosyan S, Shibata D, Perucho M. Ubiquitous somatic mutations in simple repeated sequences reveal a new mechanism for colonic carcinogenesis. *Nature* 1993;**363**:558-61.
- 6 **Thibodeau SN**, Bren G, Schaid D. Microsatellite instability in cancer of the proximal colon. *Science* 1993;**260**:816-19.
- 7 **Markowitz S**, Wang J, Myeroff L, Parsons R, Sun L, Lutterbaugh J, Fan RS, Zborowska E, Kinzler KW, Vogelstein B, Brattain M, Willson JKV. Inactivation of the type II TGF- β receptor in colon cancer cells with microsatellite instability. *Science* 1995;**268**:1336-8.
- 8 **Souza RF**, Appel R, Yin J, Wang S, Smolinski KN, Abraham JM, Zou TT, Shi YQ, Lei J, Cottrell J, Cymes K, Biden K, Simms L, Leggett B, Lynch PM, Frazier M, Powell SM, Harpaz N, Sugimura H, Young J, Meltzer SJ. Microsatellite instability in the insulin-like growth factor II receptor gene in gastrointestinal tumours. *Nat Genet* 1996;**14**:255-7.
- 9 **Rampino N**, Yamamoto H, Ionov Y, Li Y, Sawai H, Reed JC, Perucho M. Somatic frameshift mutations in the BAX gene in colon cancers of the microsatellite mutator phenotype. *Science* 1997;**275**:967-9.
- 10 **Malkhosyan S**, Rampino N, Yamamoto H, Perucho M. Frameshift mutator mutations. *Nature* 1996;**382**:499-500.
- 11 **Hornstein OP**, Knickenberg M. Perifollicular fibromatosis cutis with polyps of the colon - a cutaneo-intestinal syndrome sui generis. *Arch Dermatol Res* 1975;**253**:161-75.
- 12 **Le Guyadec Y**, Dufau JP, Poulain JF, Vaylet F, Frossin M, Lanternier G. Trichodiscomes multiples associes a une polypose colique. *Ann Dermatol Venereol* 1998;**125**:717-19.
- 13 **Schachtsschabel AA**, Kuster W, Happle R. Perifollikulare fibrome der haut und kolonpolyphen: Hornstein-Knickenberg syndrom. *Hautarzt* 1996;**47**:304-6.
- 14 **Zbar B**, Alvord WG, Glenn G, Turner M, Pavlovich CP, Schmidt L, Walther M, Choyke P, Weirich G, Hewitt SM, Duray P, Gabril F, Greenberg C, Merino MJ, Toro J, Linehan WM. Risk of renal and colonic neoplasms and spontaneous pneumothorax in the Birt-Hogg-Dube syndrome. *Cancer Epidemiol Biomarkers Prev* 2002;**11**:393-400.
- 15 **Khoo SK**, Giraud S, Kahnoski K, Chen J, Motorna O, Nickolov R, Binet O, Lambert D, Friedel J, Levy R, Ferlicot S, Wolkenstein P, Hammel P, Bergerheim U, Hedblad MA, Bradley M, The BT, Nordenskjold M, Richard S. Clinical and genetic studies of Birt-Hogg-Dube syndrome. *J Med Genet* 2002;**39**:906-12.
- 16 **Shin KH**, Park YJ, Park JG. PTEN gene mutations in colorectal cancers displaying microsatellite instability. *Cancer Lett* 2001;**174**:189-94.
- 17 **Ku JL**, Yoon KA, Kim DY, Park JG. Mutations in hMSH6 alone are not sufficient to cause the microsatellite instability in colorectal cancer cell lines. *Eur J Cancer* 1999;**35**:1724-9.
- 18 **Lengauer C**, Kinzler KW, Vogelstein B. DNA methylation and genetic instability in colorectal cancer cells. *Proc Natl Acad Sci USA* 1997;**94**:2545-50.
- 19 **Park JG**, Oie HK, Sugarbaker PH, Henslee JG, Chen TR, Johnson BE, Gazdar A. Characteristics of cell lines established from human colorectal carcinoma. *Cancer Res* 1987;**47**:6710-18.
- 20 **Duval A**, Rolland S, Compoint A, Tubacher E, Iacopetta B, Thomas G, Hamelin R. Evolution of instability at coding and non-coding repeat sequences in human MSI-H colorectal cancers. *Hum Mol Genet* 2001;**10**:513-18.
- 21 **Duval A**, Reperant M, Hamelin R. Comparative analysis of mutation frequency of coding and non coding short mononucleotide repeats in mismatch repair deficient colorectal cancers. *Oncogene* 2002;**21**:8062-6.

Want full access but don't
have a subscription?

Pay per access

For just US\$25 you can have instant access to the whole website for 30 days. During this time you will be able to access the full text for all issues (including supplements) available. You will also be able to download and print any relevant pdf files for personal use, and take advantage of all the special features *Journal of Medical Genetics* online has to offer.

www.jmedgenet.com